



International Journal of Biopharmaceutics

Journal homepage: www.ijbonline.com

IJB

PHARMACEUTICAL FORMULATION ANALYSIS: ADVANCED ANALYTICAL METHODS IN ENSURING DRUG SAFETY, EFFICACY AND QUALITY.

Dr. Sundhararajan R¹, Raman S.G^{2*}, Afrith Ali M³, Anwar Razith A³, Bhuvrangan S³.

¹Professor and Principal, Mohamed Sathak AJ College of Pharmacy, Sholinganallur, Chennai, Tamil Nadu, 600119, India.

²Professor in-charge, Mohamed Sathak AJ College of Pharmacy, Sholinganallur, Chennai, Tamil Nadu, 600119, India.

³B.Pharm seventh semester students, Mohamed Sathak AJ College of Pharmacy, Sholinganallur, Chennai, Tamil Nadu, 600119, India.

ABSTRACT

The pharmaceutical formulation is an essential part of drug development that guarantees the safety, effectiveness, and high quality of the drug products during the entire duration they are in store. This is done through the interaction of active pharmaceutical ingredients (APIs) and excipients, although stability, bioavailability and release control are emphasized. Formulation analysis helps to investigate the physical, chemical and biological characteristics of drug products to determine their functionality under clinical circumstances. The rates and consistency of drug release is measured by using various analytical approaches, including dissolution testing and particle size analysis as well as by conducting studies on stability. It is also analyzed in the manufacturing process in which there is uniformity in various production batches. Moreover, the modern formulation analysis takes into consideration the pharmacodynamics and pharmacokinetics to maximize drug dosage and therapeutic effects. As the pharmaceutical industry changes, formulation analysis is also more complicated and the use of high-techniques, such as HPLC and mass spectrometry, are necessary in order to face such issues as ingredient compatibility and compliance with regulations.

Key words: Pharmaceutical Formulation, Drug Analysis, Stability Testing, Bioavailability, Analytical Methods.

INTRODUCTION

Pharmaceutical formulation analysis is one of the basic areas in the pharmaceutical sciences, as it aims at the design, creation and testing of drug products so as to determine the safety, efficacy and quality of the drug products. Formulation of pharmaceutical products entails the interaction of active pharmaceutical ingredients (APIs), and excipients to formulate an appropriate dose. The following formulations are produced to have the drug delivered in a safe and effective way and the desired therapeutic effect.

Corresponding Author

Raman SG

E Mail ID: rtanush2015@gmail.com

Formulation analysis is used to examine physical, chemical, and biological properties of the drug product, to ascertain that it is of the required standards to use in clinical practice. The subject is vital in terms of drug development, as well as, the regulatory approval aspect, since it assures that the pharmaceutical products are safe, stable and effective throughout their shelf life. Pharmaceutical formulation begins with the identification and development of the API after which the compound is mixed with other excipients like binders, fillers, preservatives and stabilizers. [1] These excipients are significant in maintaining the stability, bioavailability, as well as the controlled release of the drug. Tablets, capsules, oral liquids, creams, ointments, and injectables may be available in different formulations thus they do not

use the same methods to analyze them and determine their physical and chemical characteristics such as dissolution rate, stability, and bioavailability. These are essential in order to find out how the drug will perform in the clinical conditions whether the drug will perform as it is supposed in the body. The formulation analysis is also the main element in determining the manufacturing procedure and scaling-up of production. The consistency of the batches is necessary to ensure the quality of the drugs, and the formulation analysis will give information on the variability possibilities in the production. This is whereby instrumental and non-instrumental methods are employed, i.e. chromatography, spectroscopy and microscopy to analyze different properties of the drug formulation. The tests would be imperative in making modifications to the formulation process and to verify that the product is up to the required standard to be used by patients. [2] Also, the current pharmaceutical formulation analysis uses the knowledge of the pharmacokinetics and pharmacodynamics of the drug. The pharmacokinetics entails the absorption, distribution, metabolism, and excretion of the drug and assists in establishment of the most suitable dosage and regimen of the treatment. Pharmacodynamics on the other hand studies the effects of the drug on the body and it gives details about the interaction of the drug with the receptors and the targets in the body. The aforementioned factors play a critical role in assessing the efficacy and safety of a drug formulation and guaranteeing the therapeutic response. The pharmaceutical industry has had a constant evolution, and formulation analysis has grown more complicated particularly with the introduction of new drug delivery mechanisms, combination therapeutic regimens, and biologics. [3] The difficulties of the formulation analysis including interactions between formulation ingredients, the development of novel analytical methods, and guarantees of the compliance with the regulations have made it a rather sophisticated discipline. Consequently, pharmaceutical researchers have to use top levels of analysis in ensuring that drugs are safeguarded and effective. This makes sure that new drug products are of the top standards of quality and best therapeutic effect on patients. With the pharmaceutical industry ever inventive, formulation analysis is a very important aspect when it comes to the development of new and better drug therapies.

The book presents an introduction to Pharmaceutical Formulation Analysis.

The study of pharmaceutical formulation is an important field of pharmaceutical sciences, which is concerned with drug product designing and evaluation of their physical and chemical characteristics. Formulation analysis aims at making sure that a drug product is of the highest quality, safe, and effective. Pharmaceutical formulations are a combination of active pharmaceutical ingredients (APIs) and other excipients, which facilitates drug delivery, stability and efficacy. [4] Formulation analysis plays a central role in ensuring that there is a correct interaction between these components, efficacy of drugs and maximizing stability and release profile of drugs.

To prepare the pharmaceutical formulation process, it starts with the identification and development of the active compound that is then subjected to several excipients like binders, fillers, preservatives, and stabilizers to form the end dosage form. Tablets, capsules, oral liquids, creams, ointments and injectables are some of the different formulations. In every formulation, there are distinct analysis methods to determine the physical and chemical properties of the formulations including rate of dissolution, stability, and bioavailability. Such assessments play a vital role in regulatory clearance and also in establishing the appropriateness of a formulation to use in the clinical setting.

Pharmaceutical formulation analysis does not just examine the composition and stability of the drug but also examines the manufacturing process and the up-scaling of production processes. This is essential so that there is consistency between various production batches. The analysis is conducted in the form of instrumental and non-instrumental analysis in the form of chromatography, spectroscopy, and microscopy. [5] The findings assist drug designers to make the required changes to formulae, and the product will be of the expected standards to be used and distributed in the clinical setting.

Further, the contemporary study of pharmaceutical formulation is also encompassed with an insight of the pharmacokinetics and pharmacodynamics of the drug, the pharmacological pathway of how the drug is absorbed, disseminated, transformed, and expelled by the body. The aspects are crucial in deciding acceptable doses and regimens of treatment. With the constantly changing industry, the analysis of the formulation of pharmaceuticals is becoming more complicated, such that the medications offered are safe and effective to the patients.



Figure1: Overview of Pharmaceutical Formulation Analysis:

The important methods of analysis used in formulation testing.

The testing of the pharmaceutical formulation is based on a number of important analytical methods that help to guarantee the drug product is safe, effective and of high quality. These are necessary techniques in describing the physical, chemical, and biological characteristics of the formulation.[6] The ability to measure various aspects, such as stability of drug, bioavailability, release profile, and overall therapeutic effect is measured analytically. These analytical techniques also allow pharmaceutical scientists to streamline formulations and their own

performance over time to ensure that the drug product is acceptable in terms of regulatory standards and that they produce the desired therapeutic effect. Major analytical techniques are that of particle size and surface area, dissolution, and release profile. All of these approaches are valuable in terms of information provided about the nature of the formulation and finally leads to the generation of safe and effective drugs. With the implementation of these techniques, the manufacturers can determine how the drug will perform within the human body and as such, consistency is guaranteed in its behavior within the different batches. [7]

Table 1: Common Pharmaceutical Formulations and Their Analytical Methods

Formulation Type	Analytical Methods	Key Considerations
Tablets	Dissolution testing, content uniformity	Drug release rate, consistency
Capsules	Hardness testing, dissolution testing	Uniformity, bioavailability
Oral Liquids	Stability testing, pH, viscosity testing	Homogeneity, shelf-life
Injectables	Osmolality, pH, stability testing	Compatibility with physiological conditions
Creams/Ointments	Viscosity, stability, release profile	Consistency,

The particle size and surface area analysis were conducted to provide the overall size and surface area of a sample after the experiment.

The analysis of particle size and surface area plays an important role in assessing the performance of pharmaceutical formulations particularly to the solid forms of dosages, such as tablets, powders, and suspensions. Drug particles dimension and surface area can influence greatly their solubility, bioavailability and general effectiveness as therapeutic agents. [8] The smaller the particles, the higher the surface area and thus absorption of drugs that are poorly soluble in the body can be improved. Particle size distribution and surface area is usually measured by using techniques like laser diffraction, dynamic light scattering (DLS), and scanning electron

microscopy (SEM). Laser diffraction is extensively utilized due to the fact that it can be used to measure a very wide size of the particle, whereas DLS can be applied to study nanoparticles or suspended particles. The techniques assist in identifying the optimal size of the particle and surface area that can be used to achieve a specific drug release profile and therapeutic effect. [9] Also, the analysis of particle size is essential to provide consistency in the various batches of the drug product since the difference in particle size may result in alterations in the drug performance. This analytical method is also useful in optimization of formulation approaches so as to increase drug efficacy and stability.

Table 2: Particle Size and Surface Area Techniques

Technique	Description	Application	Advantages
Laser Diffraction	Measures particle size distribution via laser light scattering	Suitable for a wide range of particle sizes	High-speed, broad range measurement
Dynamic Light Scattering (DLS)	Measures light scattering patterns from particles in suspension	Ideal for nanoparticles and colloidal systems	Can measure very small particles (1-1000 nm)
Scanning Electron Microscopy (SEM)	Provides high-resolution imaging of particle surface	Visualization of particle morphology	High precision and detailed surface analysis

Dissolution Testing and Release Profile Investigations.

Dissolution testing and release profile studies are the basic methods of testing the release rate of drug in the dosage form and availability to the body. Dissolution testing is used to determine the rate and degree of drug dissolution in a simulated gastrointestinal fluid that is important in determining the bioavailability of the drug. This is of particular significance to oral solid dosage forms of tablets and capsules where the medication has to be dissolved in the stomach or intestines before it can be absorbed.[10] The outcomes of the dissolution testing give useful data on the performance of the formulation regarding the release kinetics and the capacity of the formulation to perform the necessary therapeutic standard.

Different equipment is applied in dissolution tests, including the USP Apparatus 1 (basket method) and 2 (paddle method) that enables the testing conditions to be carefully controlled like temperature, agitation and pH of the dissolution medium. Release profile studies are an extra step, as they analyze the release of the drug with time, which gives a complete picture of the release and absorption of the drug into the bloodstream. Such studies play a critical role in determining the therapeutic effect of the drug since they are used in optimizing the rate at which the formulation is released and to ensure that the drug produces the desired therapeutic effect in the required duration.

solid dosage forms Analytical methods.

Pharmaceutical formulation process entails development and testing of solid dosage, including pills and capsules. The analytical techniques which are applied to solid dosage forms are aimed at measuring the different physical, chemical and mechanical characteristics which determine the performance of the drug. Such techniques are used to ascertain the uniformity of the drug content, its stability, solubility, as well as its quality. Solid dosage form consistency and reliability is very important because any difference can have a severe effect on the bioavailability and the therapeutic effect of the drug[11]. Other methods of analysis are used as well as the tablet and capsule analysis, powder flow/ compressibility test. Such tests give useful information of the behavior of the drug in various circumstances in production and after administration, in order to optimize the formulation with regard to consistency in manufacturing and therapeutic efficacy.

Tablet and Capsule Analysis

Tablet and capsule analysis are especially important in assessing the quality of solid dosage form, safety, and efficacy. In the analysis, various characteristics, which include uniformity of content, hardness, disintegration time, and the dissolution characteristics are examined. The analysis of a tablet and capsule, the measurement of content uniformity is one of the most widely used tests that provides guarantees that the required amount of active pharmaceutical ingredient (API) is in each unit of dosage.[12] The uniformity of the content is very vital in providing the patient with the right amount of the drug. The next significant parameters are the hardness and friability tests that can be used to determine the resistance to breakage and the mechanical strength in handling, transportation, and storage. A tablet or capsule that is excessively soft can fracture or crumble hence making dosing inconsistent whereas too hard may influence the speed at which the drug is dissolved. The disintegration test determines the duration in which the tablet falls apart into small particles in the gastrointestinal tract to release the drug and allow its absorption at the right time. Dissolution testing that determines the rate and degree of drug release out of the pill or capsule is important in the determination of the bioavailability of the drug and its therapeutic competence. These tests aid in optimization of the formulation to make sure that the drug works in a consistent manner under the physiological conditions. Tablet and capsule analysis is necessary to be able to approve the drug by the regulator, each batch of which should meet the necessary quality standards and effectively reach the patient.

1.3.2 Powder Flow and Compressibility Test.

Powder flow and compressibility tests are necessary in assessing the physical characteristics of the

raw material to be used in solid dosage forms and the properties of the powder during the manufacturing of the product like compression of the powder into tablets and capsule filling. [13] These tests determine the flowability of a powder and compressibility into a tablet or a capsule. The free-flowing nature of the powder is important in allowing consistency in the manufacturing process, especially in a high-speed rectangular press, or capsule fillers, when the poor flow results in inconsistency in the filling and dosing. Angle of repose, Hausner ratio, and Carr index are some of the most common tests used to determine the properties of powder flow. Angle of repose offers the index of the capacity of the powder to produce a pile whereby a high angle means low flowability. Hausner ratio and the index of Carr are used to evaluate the correlation between bulk density and tapped density of powder, which provides information about the compressibility of the powder and the tendency to produce fine grains in the course of formulation. Compressibility testing determines how the powder can be pressed into solid tablets which gives information on how the powder will be able to make hard enough tablets that do not have high friability. [14] Such tests are used to maximise the formulation to choose powders with good flow, compressibility, and to form tablets or capsules that have the right mechanical properties. Also, the compressibility testing is able to determine possible problems, like capping, lamination, or a poor build-up of the pill, which might influence the quality and performance of the product.

Analytical methods of Liquid dosage form.

The most common liquid dosage form is solutions, suspensions, syrups, and emulsions, which are used because of their ease of administration, especially in the delivery of pediatric, geriatric, and parenteral drugs. Liquid dosage forms require analytical methods that are critical to guarantee their stability, efficacy, and safety. Such techniques will assist in determining the physical, chemical, and microbiological characteristics of the formulation to make it of the necessary quality to be used by the patients. [15] The important liquid formulations tests are stability tests, viscosity tests, pH and osmolality tests. Such analytical methods give useful data about the behavior of the formulation given storage condition, flow properties and suitability to be administered to the patients. Stability and viscosity tests are used to ensure the drug retains the desired potency and consistency due to passage of time whereas, pH and osmolality tests are used to ensure the liquid formulation is not incompatible with the physiological state of the body, which would result in irritation or adverse reactions.



Figure2: Analytical Methods for Solid Dosage Forms

Table 3: Common Analytical Methods for Liquid Dosage Forms

Test Type	Technique Used	Purpose	Example Application
Stability Testing	Accelerated stability testing	Evaluate long-term storage stability	Predicting shelf-life of oral liquids
Viscosity Testing	Viscometer, Rheometer	Measure flow properties	Ensuring ease of administration in suspensions
pH Testing	pH Meter	Ensure formulation compatibility with body conditions	Oral and intravenous drug formulations
Osmolality Testing	Osmometer	Ensure	Parenteral drug products

The measurements of stability and viscosity are presented.

The parameters of stability and viscosity used in the context of liquid dosages are needed to guarantee the quality of products, their potency, and safety during their shelf life. Stability testing assesses a drug formulation in terms of its ability to retain its physical, chemical and microbiological characteristics over time in various environmental factors, including temperature, humidity and light, etc. This plays an important role in establishing the necessary expiration date of the drug and monitoring the fact that the drug will be rather efficient and safe to use till this moment. Other stability tests include accelerated stability tests whereby the samples are put under extreme conditions to determine their stability in the long run. On the other hand, the measurement of viscosity evaluates the flow characteristics of the liquid formulation which is a critical aspect to the adequate administration of the drug. As an example, excessive viscosity of liquid formulations can make them hard to swallow or to inject, and a low viscosity may fail to offer the intended sustained-release properties. [16] A viscometer or rheometer can be used to measure viscosity and the results of this test can be used to optimize the formulation to be easy to administer and predictable in its therapeutic effect. Moreover, viscosity is significant to homogeneity of suspensions and emulsions, making sure that the active ingredient is evenly spread all through the liquid. The stability and viscosity tests are also critical to the success of the development of liquid dosage forms, as it guarantees that the formulation is retaining its intended therapeutic effects, is safe to the patient, and is readily administered.

pH and Osmolality Testing

pH and osmolality assays play a vital role in determining compatibility and safety of liquid dosage formulations, especially in oral, intravenous, or ocular administration. pH testing determines acidity or alkalinity of a liquid formulation that may greatly affect solubility, stability and bioavailability of the active pharmaceutical ingredient (API). Most drugs have got a fixed pH within which they are stable and outside this range then they might either be degraded or rendered ineffective. As an example, any formulation that is to be taken as an oral agent must have a compatible pH that does not only accommodate to the stability of the drug but also the gastrointestinal conditions of the patient. Equally, intravenous (IV) formulations should maintain a pH near to the physiological pH (approximately 7.4) in order to prevent irritation or harm to blood vessels when it is administered, keeping the formulation within the desired pH range to achieve the desired therapeutic effect. Osmolality testing is used to determine the concentration of solute particles in a liquid formulation that is of special concern to parenteral or intravenous products. [17,18] Osmolality plays an important role in determining the

osmotic pressure of a solution and irregularities with that of physiological levels may result in cell damage or undesirable reactions. Osmolality testing of formulations will determine the isotonicity with body fluids and minimize chances of irritation or hemolysis in the case of intravenous administration. An osmometer can be used to measure osmolality, which is necessary to preserve the safety and effectiveness of intravenous, ophthalmic or other specialized liquid preparations. The pH and osmolality assay are both invaluable in the formulation and quality assurance of liquid dosage forms, meaning the formulation is safe, effective and compatible with the physiological factors of the body.[19,20]

Dilemmas of Formulation Analysis.

Although formulation analysis is a very necessary field to maintain the quality, safety and efficacy of the drug, it poses many challenges that cannot be overlooked and handled without a lot of care and expertise. Modern pharmaceutical formulations, which usually require a mixture of several active ingredients and excipients, require an in-depth understanding of the interactions between many different components and the capacity to substantiate complex formulations. [21,22] These difficulties are enhanced by the variety of different types of formulations (e.g., oral, injectable, topical) and the spectrum of physicochemical characteristics, which should be considered. In efficient formulation analysis, issues of ingredient compatibility, ingredient stability and analytical method reproducibility should be addressed. Addressing these issues is essential to the formation of pharmaceutical products that are not only of a standard that is paralleled by the regulatory authorities but that also would give the same therapeutic effect on diverse populations of patients. Some of the most important issues in the study of formulation are the interactions of the ingredients of a formulations and the analytical validation of complex formulations. [23]

Interaction between Formulation Ingredients.

The interactions of the different formulation ingredients are one of the main issues in the pharmaceutical formulation analysis. The presence of numerous excipients (binders, preservatives, stabilizers, etc.) in addition to the active pharmaceutical ingredient (API) may result in unplanned interactions that interfere with the stability, bioavailability and, overall, functionality of the drug. These interactions may occur in various forms, including degradation of active ingredient, alteration of dissolution profile of the active ingredient, or the generation of unwanted by-products. [24,25] As an illustration, solubility of the API can be changed by some excipients or, the excipients can change the pH of the formulation, which may affect the stability or efficacy of the drug. Also, the excipients such as surfactants and polymers may modify the rate of the release of the drug

which could cause interference with the bioavailability and therapeutic activity of a drug. In order to overcome these challenges, researchers apply the methods of stability, pre-formulation and compatibility studies to determine the interactions between the ingredients prior to formulation. The possible adverse reactions that may occur between the API and excipients need to be detected and avoided at all costs so that the end product can be as effective as desired. [26] The design of formulating properly and the selection of excipients is very important in the prevention of unwanted interactions that might lead to the undermining of safety and efficacy of the drug.

Analytical Check of Complex Formulations.

Another major problem of pharmaceutical formulation analysis is the analysis of complex formulations; whose validation is analytical. Multicomponent formulations are frequently a mixture of several APIs and excipients which have physical, chemical and biological characteristics. This is a complex issue and thus necessitates the application of advanced analytical methods in a manner that will enhance the formulation attaining the desired quality threshold and thus be effective.[27] Analytical methods should be validated in order to make sure that methods in the analysis of the formulation are accurate, precise, specific, and strong.

When complex formulations, particularly those containing controlled-release systems, biopharmaceutical drug delivery systems, or combination therapies are to be developed, it is imperative to devise and test protocols that have the potential to reliably determine the release profiles, stability, and bioavailability of the drug.[28] This may need sophisticated methods like high-performance liquid chromatography (HPLC), mass spectrometry (MS) and spectrophotometry, in addition to dedicated methods of measuring particle size, dissolution, and stability. The other aspect in analytical validation is whether the testing methods are appropriate to the particular formulation and purpose of use. This may especially be a problem with novel or combination therapy formulations, whereby there may be few standards by which to provide a reference standard, and the new, innovative validation protocols must be formulated [29]. The validation should be rigorous to the extent that the analytical procedures should have the competency of measuring the quality and performance of the complex formulations over their shelf life, both in terms of regulatory compliance and patient safety. Superb validation of methods of analysis also assists in the consistency and reliability of drug products at various manufacturing batches. [30]



Figure2: Challenges in Pharmaceutical Formulation Analysis

Conclusion

The analysis of pharmaceutical formulations is very important in the design, development, and regulation of drug products to ensure that they are safe, effective and of quality during their shelf life. It starts with the identification of active pharmaceutical ingredients (APIs), then the addition of excipients (binders, fillers, preservatives, and stabilizers) which are necessary to stabilize, sustain bioavailability, and release of drugs. The purpose of the formulation analysis is to make sure that the association between the APIs and excipients is ideal, which will help to create the intended therapeutic effect and avoid any adverse reaction.

The identification of physical, chemical and biological properties of the drug product is one of the most important elements of pharmaceutical formulation analysis. Dissolution test, particle size, and stability tests are analytical techniques that are used to quantify the rate of release of the drug, its ability to be dissolved, and its consistency. Such procedures are essential in estimating the performance of the drug in human body given the clinical conditions. Moreover, regulatory bodies demand such analyses to ensure that the formulation does not violate the formulation standards so that the drug is safe and effective to be used by the patients. The analysis does not end at the evaluation of the composition and stability of the drug. It goes further to assess the manufacturing process whereby every lot of the drug product is consistent and re-producible. The drug formulation is analyzed using instrumental techniques (chromatography, spectroscopy, and microscopy) and non-instrumental techniques in order to identify any potential problems. These analyses inspire

important information providing directions to the formulation process, which optimize the performance and quality of the drug. Also, the modern study of pharmaceutical formulation includes knowledge of pharmacodynamics and pharmacokinetics. Pharmacokinetics which concerns itself with the absorption, distribution, metabolism and excretion of the drug is critical to the determination of proper dosage regimens. Pharmacodynamics, on the other hand, is the study of the drug and its interaction with the body and its activity on different biological pathways. These factors combined are the basis on which the therapeutic efficacy and safety of the drug can be evaluated. The pharmaceutical industry keeps changing as the new drug delivery system, biologics, and combination therapies are created, and so, the formulation analysis becomes more complicated. Such challenges as the compatibility of the ingredients, the stability of the formulations, and the analysis validation are to be considered carefully in order to guarantee the intended results. These obstacles necessitate sophisticated analytical techniques, including the high-performance liquid chromatography (HPLC), mass spectrometry (MS), and UV spectroscopy to ascertain the dependability and the performance of complicated formulations. To sum up, pharmaceutical formulation analysis is a complex field, which is essential in the formulation of safe and effective drug treatments. As the pharmaceutical science constantly develops, the formulation analysis will also keep progressing, responding to the arising issues and guaranteeing the highest levels of safety, effectiveness, and quality of the drug to the patient.

REFERENCES

1. Bakhrushina EO, Shumkova MM, Avdonina YV, Ananian AA, Babazadeh M, Pouya G, *et al.* Transdermal Drug Delivery Systems: Methods for Enhancing Skin Permeability and Their Evaluation. *Pharmaceutics* 17, 2025, 936.
2. Dhull P, Dunuweera S, Bietsch J, Bandu R, Wannere C, Achanta S, *et al.* Recent advances and application of liquid chromatography in pharmaceutical industry. *Journal of Liquid Chromatography & Related Technologies* 48, 2025, 168–187.
3. Comoglu T, & Ozyilmaz ED. Pharmaceutical excipients in pediatric and geriatric drug formulations: safety, efficacy, and regulatory perspectives. *Pharmaceutical Development and Technology* 1–9, 2024.
4. Bunaciu AA, Hoang VD, Aboul-Enein HY. *et al.* Fourier transform infrared spectroscopy used in drug excipients compatibility studies. *Applied Spectroscopy Reviews* 60, 2024, 385–403.
5. Chen Y, Tang Y, Li Y, Rui Y, Zhang P. *et al.* Enhancing the Efficacy of Active Pharmaceutical Ingredients in Medicinal Plants through Nanoformulations: A Promising Field. *Nanomaterials* 14, 2024, 1598.
6. Finotti Cordeiro C, Lopardi Franco L, Teixeira Carvalho D, Bonfilio R. *et al.* Impurities in Active Pharmaceutical Ingredients and Drug Products: A Critical Review. *Critical Reviews in Analytical Chemistry* 56, 2024, 1–21.
7. Das A, & Prajapati P. Navigating pharmaceuticals: microfluidic devices in analytical and formulation sciences. *Discov Chem* 2, 2025.
8. Kapoor DU, Kukkar RR, Gaur M, Prajapati BG, Malviya R. *et al.* The Potential of Quality Target Product Profile in the Optimization of Nanoemulsions. *CNANOM* 15, 2025, 4–17.
9. Kovács B, Tőkés E-O, Kelemen ÉK, Zöldi K, Boda F, Suba E, *et al.* The Effect of Formulation Variables on the Manufacturability of Clopidogrel Tablets via Fluidized Hot-Melt Granulation-From the Lab Scale to the Pilot Scale. *Pharmaceutics* 16, 2024, 391.

10. Ahmed Y, Mahmoud AAK, Ludasi K, Sovány T. *et al.* Advances in Loading Techniques and Quality by Design for Fused Deposition Modeling in Pharmaceutical Production: A Systematic Review. *Pharmaceutics* 17, 2024,1496.
11. Chen Z, Lian JZ, Zhu H, Zhang J, Zhang Y, Xiang X, *et al.* Application of Life Cycle Assessment in the pharmaceutical industry: A critical review. *Journal of Cleaner Production* 459, 2024, 142550.
12. Alikhani MS, Nazari M, Hatamkhani S. Enhancing antibiotic therapy through comprehensive pharmacokinetic/pharmacodynamic principles. *Front Cell Infect Microbiol* 15, 2025.
13. Liu J, Nordin JZ, McLachlan AJ, Chrzanowski W. *et al.* Extracellular vesicles as the next-generation modulators of pharmacokinetics and pharmacodynamics of medications and their potential as adjuvant therapeutics. *Clinical & Translational Med* 14, 2024.
14. Garcia-Cremades M, Solans BP, Strydom N, Vrijens B, Pillai GC, Shaffer C, *et al.* Emerging Therapeutics, Technologies, and Drug Development Strategies to Address Patient Nonadherence and Improve Tuberculosis Treatment. *Annu Rev Pharmacol Toxicol* 62, 2021, 197–210.
15. Zhu T, Liu B, Chen N, Liu Y, Wang Z, Tian X. *et al.* Artificial Intelligence-Driven Innovations in Pharmaceutical Development and Drug Delivery Systems. *CTMC* 25, 2025,2937–2951.
16. B S, & Ghosh A. Mechanistic Insights into Amorphous Solid Dispersions: Bridging Theory and Practice in Drug Delivery. *Pharm Res* 42, 2025,1–23.
17. Chaudhary A, & Shambhakar S. Nanotechnology in Drug Delivery: Overcoming Poor Solubility Challenges through Nanoformulations. *CNANOM* 14, 2024,200–211.
18. Shin Y, Kim M, Kim C, Jeon H, Koo J, Oh J, *et al.* Development and Characterization of Olaparib-Loaded Solid Self-Nanoemulsifying Drug Delivery System (S-SNEDDS) for Pharmaceutical Applications. *AAPS PharmSciTech* 25,2024.
19. Shrivastava A, Mandal S, Pattanayek SK, Rathore AS. *et al.* Rapid Estimation of Size-Based Heterogeneity in Monoclonal Antibodies by Machine Learning-Enhanced Dynamic Light Scattering. *Anal Chem* 95, 2023,8299–8309.
20. Zahariev N, Draganova M, Zagorchev P, Pilicheva B. *et al.* Casein-Based Nanoparticles: A Potential Tool for the Delivery of Daunorubicin in Acute Lymphocytic Leukemia. *Pharmaceutics* 15, 2023,471.
21. Jayachandran P, Ilango S, Suseela V, Nirmaladevi R, Shaik MR, Khan M, *et al.* Green Synthesized Silver Nanoparticle-Loaded Liposome-Based Nanoarchitectonics for Cancer Management: In Vitro Drug Release Analysis. *Biomedicines* 11, 2023,217.
22. Olsson M, Storm R, Björn L, Lilja V, Krupnik L, Chen Y, *et al.* Phase-separated polymer blends for controlled drug delivery by tuning morphology. *Commun Mater* 5, 2024.
23. Van Haaren C, De Bock M, Kazarian SG. *et al.* Advances in ATR-FTIR Spectroscopic Imaging for the Analysis of Tablet Dissolution and Drug Release. *Molecules* 28, 2023,4705.
24. Rocha B, De Morais LA, Viana MC, Carneiro G. *et al.* Promising strategies for improving oral bioavailability of poor water-soluble drugs. *Expert Opinion on Drug Discovery*;18, 2023,615–627.
25. Singh P, Sharma S, Sharma PK, Alam A. Drug Polymorphism: An Important Pre-formulation Tool in the Formulation Development of a Dosage Form. *CPC* 14, 2024,2–19.
26. Tanga S, Ramburrun P, Aucamp M. From Liquid SNEDDS to Solid SNEDDS: A Comprehensive Review of Their Development and Pharmaceutical Applications. *AAPS J* 28, 2025.
27. Preeti P, Sambhakar S, Malik R, Bhatia S, Al Harrasi A, Rani C, *et al.* Nanoemulsion: An Emerging Novel Technology for Improving the Bioavailability of Drugs. *Scientifica* 2023:1–25.
28. Mohite P, Singh S, Pawar A, Sangale A, Prajapati BG. Lipid-based oral formulation in capsules to improve the delivery of poorly water-soluble drugs. *Front Drug Deliv* 3, 2023.
29. Janssen PHM, Fathollahi S, Bekaert B, Vanderroost D, Roelofs T, Vanhoorne V, *et al.* Impact of material properties and process parameters on tablet quality in a continuous direct compression line. *Powder Technology* 424, 2023,118520.
30. Tikhomirov E, Åhlén M, Di Gallo N, Strømme M, Kipping T, Quodbach J, *et al.* Selective laser sintering additive manufacturing of dosage forms: Effect of powder formulation and process parameters on the physical properties of printed tablets. *International Journal of Pharmaceutics* 635, 2023,122780.