



International Journal of Biopharmaceutics

Journal homepage: www.ijbonline.com

IJB

MODERN APPROACHES IN PHARMACEUTICAL ANALYSIS: VIBRATIONAL CHEMICAL IMAGING AND CHEMOMETRIC TECHNIQUES

Dr. Bhavani J*

Vice Principal cum HOD, Department of Pharmaceutics, St.Mary's Institute Of Pharmacy, Chinthamani, Puliangudi. -
627855, Tamilnadu, India.

ABSTRACT

Modern pharmaceutical analysis has undergone a significant transformation with the advent of vibrational chemical imaging (VCI) integrated with chemo metric techniques, enabling comprehensive, non-destructive, and high-resolution characterization of complex drug systems. VCI, employing spectroscopic modalities such as Fourier Transform Infrared (FTIR), Raman, and Near-Infrared (NIR) imaging, facilitates simultaneous acquisition of spatial and molecular information, thereby overcoming the limitations of conventional analytical methods. These techniques allow detailed visualization of active pharmaceutical ingredient (API) distribution, polymorphism, crystallinity, and drug–excipient interactions within pharmaceutical formulations. However, the large and complex datasets generated through hyperspectral imaging necessitate the use of chemo metric tools for efficient data processing and interpretation. Chemo metric methods, including principal component analysis (PCA), partial least squares (PLS), multivariate curve resolution (MCR), and machine learning approaches, play a crucial role in data pre-processing, pattern recognition, classification, and quantitative analysis. The integration of VCI and chemometrics has significantly enhanced pharmaceutical applications such as solid dosage form characterization, quality control, and process monitoring. Furthermore, these approaches align with regulatory frameworks such as Process Analytical Technology (PAT) and Quality by Design (QbD), supporting real-time monitoring and improved process understanding. Advances in imaging resolution, detector technologies, and data integration strategies have further strengthened the applicability of these techniques in pharmaceutical research and manufacturing. Despite challenges related to instrumentation cost, data complexity, and standardization, ongoing developments in automation and artificial intelligence are expected to drive wider adoption. Overall, VCI coupled with chemometric analysis represents a powerful and evolving analytical paradigm that supports the development of safe, effective, and high-quality pharmaceutical products.

Key words: Vibrational Chemical Imaging; Chemo metrics; Pharmaceutical Analysis; Spectroscopic Imaging.

INTRODUCTION

Vibrational chemical imaging (VCI) is now recognized as an effective method of analysis in the pharmaceutical sciences, allowing the determination of spatial and chemical data of complex drug systems simultaneously

Corresponding Author

Dr. Bhavani J

VCI is able to characterize pharmaceutical materials at the micro- and nanoscale by non-destructively using the vibrational spectroscopy methodologies, including infrared (IR) and Raman spectroscopy, with state-of-the-art imaging capabilities. The method gives informative details of the molecular composition, distribution, and interactions of active pharmaceutical ingredients (APIs) and excipients in solid dosage forms, which are important compared to other analytical approaches which are

mainly devoid of spatial resolution. (Bakhrushina EO *et al.*, 2025) Nowadays, the growing complexity of drug delivery systems, such as the multi-component systems and controlled-release matrices, as well as nanostructured drug delivery systems, has led to the need to use an increasingly complex analysis method like VCI. Nevertheless, the quantity of spectral information produced by these imaging modalities is a significant challenge in data interpretation and extraction of valuable information. Here is where the chemometrics, the use of mathematical and statistical techniques in chemical data, is important. The chemometric methods are used in preprocessing, modeling, and interpretation of huge multidimensional data to allow the proper qualitative and quantitative analysis of pharmaceutical systems (Bakhrushina EO *et al.*, 2022). Principal component analysis (PCA), partial least squares (PLS), cluster analysis, and multivariate curve resolution (MCR) are commonly used in order to improve the data quality, minimize noise, find patterns, and determine correlations between spectral characteristics and chemical properties. VCI combined with chemo metric analysis has made great progress in the field of pharmaceutical research by facilitating the study of drug distribution, polymorphism, crystallinity and drug excipient compatibility, which are very important in ensuring efficacy, stability and safety of the product. Moreover, this combination method would conform to the principles of Process Analytical Technology (PAT) and Quality by Design (QbD) that aids in real-time manufacturing process monitoring and control. (A Gibbons L, *et al.*, 2021) The emphasis on the use of advanced analytical methods in the process of establishing the product quality and enabling its consistency is generally gaining popularity among regulatory agencies, which further explain the topicality of VCI and chemometrics in the present-day pharmaceutical development. These advancements notwithstanding, issues like high cost of instrumentation, complexity of data and standardization of protocols still exist. However, the process of constant improvements in imaging resolution, data-integration, and machine learning-based chemometrics is likely to improve the relevance and strength of the vibrational chemical imaging. As a result, VCI in combination with chemometrics is an innovative device in pharmaceutical analysis, providing in-depth information, which helps in the development of drugs, quality regulation, and compliance with regulatory requirements in a dynamic pharmaceutical environment. (Kim EJ, *et al.*, 2021)

Instrumentation and Imaging Modalities in Chemical Imaging

The technological base of vibrational chemical imaging (VCI) is instrumentation and imaging modalities because they allow the detailed spatial and molecular characterization of pharmaceutical systems. VCI is

mainly based on spectroscopic methods which include Fourier Transform Infrared (FTIR) imaging and Raman imaging with each having its own merits depending on how it interacts with the molecular vibration. FTIR imaging employs the use of mid-infrared radiation to investigate fundamental vibrational modes which give it strong absorption signals and high chemical specificity, which is ideal in the analysis of organic compounds, functional groups and excipient matrices within solid dosage forms. (Yang S, *et al.*, 2025) Current FTIR systems commonly have focal plane array (FPA) detectors, which allow simultaneous collection of thousands of spectra, and Chemical mapping at a high rate. Raman imaging on the other hand is a technique founded upon inelastic scattering of monochromatic light, usually a laser source, and provides excellent spatial resolution and little interference with water hence is of great benefit in the analysis of water systems, polymorphic forms, and crystalline structures. More sophisticated Raman modalities like confocal Raman microscopy also increase depth profiling and three-dimensional imaging that can be used to visualize drugs distribution in complex formulations in more detail. Along with the traditional FTIR and Raman methods, newer modalities like Near infrared (NIR) chemical imaging and coherent Raman methods including Coherent Anti- Stokes Raman Scattering (CARS) and Stimulated Raman scattering (SRS) have been developed to broaden the scope of analysis of the VCI by offering higher speed of imaging and better sensitivity. These systems are normally coupled with high-performance optics, sensitive detectors and automated sample stages in order to achieve an accurate and reproducible data acquisition. (Xie J, Gowen A, *et al.*, 2024) The instrumentation is determined by the type of sample, the spatial resolution that is needed, the depth of analysis, and sensitivity. In addition, recent developments in hyperspectral imaging have made it possible to record the complete spectral data at every pixel, producing large multidimensional data sets that not only record the spatial but also chemical diversity. These datasets together with a powerful set of chemometric methods give in-depth information on the composition and structure of pharmaceuticals. In general, the unceasing development of the instrumentation and imaging modalities has contributed greatly to the improvement of the capabilities of chemical imaging turning it into the inseparable part of pharmaceutical research, development of formulations and quality control. (ChitraRagupathy I, *et al* 2021)

Overview of Chemo metric Methods in Spectral Data Analysis

Chemometric techniques have a significant role in the analysis of spectral data produced by the vibrational chemical imaging (VCI), allowing the derivation of chemical and spatial information of a

complex and high-dimensional data. Spectral data collected by methods as FTIR, Raman, and NIR imaging usually has thousands of variables per sample, posing the problem of data redundancy, noise, and overlapping spectral components. Chemometrics is a combination of statistical, mathematical and computational analysis programs that is applied to overcome such challenges through converting raw spectral data to interpretable and actionable information. Principal component analysis (PCA) is one of the most popular chemometric methods that are unsupervised and tend to reduce the dimensionality of data and maintain the variance to help identify the patterns, cluster them, and identify the underlying tendencies in the pharmaceutical samples. (Vosough M, *et al.* 2024) Also, the partial least squares (PLS) regression is being widely used as the quantitative analysis method that enables serious correlations between the spectral data and concentration, or physicochemical characteristics of the analyte. Cluster analysis methods, such as hierarchical cluster analysis (HCA) and k-means clustering are used to cluster similar spectra and map heterogeneity of space in formulations. Moreover, multivariate curve resolution (MCR) offers the capability of breaking down complicated spectral mixtures of complex spectral mixtures into pure component spectra and concentration profiles without prior understanding of the system, thus finding particular application in multi-component pharmaceutical formulations. (Elhamdaoui O, *et al.*, 2020) Linear discriminant analysis (LDA) and support vector machines (SVM), are some of the supervised methods of classification that are useful in predictive modeling and allow to categorize correctly polymorphic form, counterfeit drugs, and quality attributes. Chemometric models are frequently combined with data preprocessing methods such as baseline correction, normalization, smoothing and derivative transformations to enhance the quality of signal and improve model performance. Chemometrics can be further applied to data reduction, model validation, outlier detection and optimization of analytical processes. As machine learning and artificial intelligence emerged, chemometric methods have further developed to be able to work with high-scale hyperspectral data, which can be manually used to make decisions in other pharmaceutical processes in real-time. In general, chemometric techniques cannot be ignored to achieve the maximum potential of spectral data in VCI, which offers greater accuracy, efficiency, and reliability in the analysis of pharmaceuticals, quality control, and drug development. (Islam MN *et al.*, 2022)

Preprocessing Techniques for Spectral Data Enhancement

The preprocessing method is a vital procedure of vibrational chemical imaging (VCI) and spectral data analysis because they improve the quality of the data and

guarantee the accurate interpretation of complicated pharmaceutical data. Raw spectral data that have been collected using methods like FTIR, Raman and NIR imaging tend to exhibit noise, baseline effects, scattering or instrumental variations that may blur valuable chemical data, and invalidate further chemometric analysis. Various preprocessing techniques are also used in order to mitigate these issues, by removing distortions and normalizing spectral profiles. Baseline correction this operation is typically employed to remove background differences due to instrument drift or fluorescence and enhance the clarity and accuracy of the peaks. (Matveeva I, *et al.*, 2022) Techniques like Savitzky-Golay filtering are used to minimize random noise, and still maintain significant spectral characteristics of data to improve signal-to-noise ratio without doing much to distort the shape of the peaks. Normalization techniques such as the use of vectors and area normalization are employed to correct sample thickness, concentration or conditions of measurements, so that there is comparability among spectra. Derivative spectroscopy and especially first and second derivatives are very common to treat the overlapping lines and enhance spectral resolution which leads to better identification of the chemical components. (Zhu Q, Liu Q, *et al.*, 2025) Multiplicative scatter correction (MSC) and standard normal variate (SNV) are scatter correction methods that are used to eliminate the impacts of light scattering and variations in particle size that are of particular concern in solid dosage forms and heterogeneous samples. Further, spectral truncation and wavelength choice aid the elimination of irrelevant or noisy areas and the analysis is condensed to informative spectral areas. Preprocessing is also important in improving the performance of chemometric models by minimizing the dimension of data and also removal of redundant information. (Khadem H, *et al.*, 2024) The preprocessing steps should be chosen and ordered with great care since improper use will result in the loss of important information or addition of artifacts. Also in the recent years, automated and adaptive preprocessing algorithms have been created which are commonly combined with machine learning models to simplify data processing and increase reproducibility. All in all, preprocessing methods are essential in converting raw spectral data into quality datasets, and thus making possible the correct analysis of qualitative and quantitative analysis in drug development, such as formulation development, quality assurance and monitoring process. (Chen Z, *et al.* 2024)

Multivariate Analysis for Pharmaceutical Applications

Multivariate analysis is of primary interest in pharmaceutical use because it can be used to interpret multivariate high-dimensional data provided by vibrational chemical imaging and alternative analysis

methods. Pharmaceutical systems are multicomponent in nature and are known to have active pharmaceutical ingredients, excipients, impurities and degradation products, which tend to generate overlapping spectral signals which cannot be separated using the univariate methods. The multivariate methods offer the tools with high power to examine such data viewing several variables at the same time and thus discovering more obscured patterns, correlations, and hidden structure of the data. (Pokhrel DR *et al.*, 2023) Principal component analysis (PCA) is a popular exploratory object reduction method when identifying clustering, outliers in a batch of medications to enable quick evaluation of changes in batch to batch variability, and product stability. To analyse quantitative data, partial least squares (PLS) regression is widely used to create calibration models that predict the changes in concentration of drug components by relying on spectral data, which is useful in testing content uniformity and developing dissolution profiles (Wang J, *et al.*, 2021). The use of linear discriminant analysis (LDA), soft independent modeling of class analogy (SIMCA), and support vector machines (SVM) on the samples are used to classify, differentiate between various polymorphic forms, and detect counterfeit or substandard drugs. Multivariate curve resolution (MCR) also allows better data interpretation of complex mixtures breaking them down into spectrums due to pure components and concentration profiles, even when no prior information is known. The methods are especially useful in the analysis of the distribution of drugs, excipient interactions, and heterogeneity in solid dosage forms. Also, Process Analytical Technology (PAT) systems incorporate multivariate techniques, which allow real-time monitoring and control of pharmaceutical manufacturers. (Dorabiala O, *et al.* 2024) They assist the Quality by Design (QbD) concepts through the provision of risk assessment, optimization of the processes, and creation of strong design space. Techniques Model validation models, such as cross-validation, and external validation are used to make sure models of multivariates are reliable and predictive in a regulatory setting. As machine learning and artificial intelligence have been integrated, multivariate analysis has also evolved to work with great volumes of data and automate the process of decision-making. In general, the multivariate analysis is used as a pillar of the contemporary pharmaceutical research and quality assurance that offers greater accuracy, efficiency, and insight into the complex drug systems. (Teng G, Wang Q, *et al.* 2021)

Application in Solid Dosage Form Characterization

Vibrational chemical imaging (VCI) coupled with chemometric analysis has turned into a vital instrument in the characterization of solid dosage forms where it offers detailed spatial and molecular data that is paramount in guaranteeing the quality, efficacy and

stability of the product. Tablets and capsules are solid dosage forms that are multicomponent complex systems that contain active pharmaceutical ingredients (APIs) and various excipients, and their homogeneous distribution and favorable interaction are required to achieve consistent therapeutic performance. The conventional methods of analysis tend to give bulk data and not to show spatial heterogeneity; VCI can be used to generate non-destructive high-resolution maps of chemical constituents in the dosage form. Such techniques as FTIR and Raman imaging can be used to visualize API distribution, determine excipient domains, and identify agglomeration or segregation problems that can influence drug release and bioavailability. Chemometric analyses are additional means through which such analyses are improved by allowing a classification, quantification and pattern identification on large spectral data sets, to ensure a valid evaluation of the homogeneity of content and the homogeneity of blends. (Kucharska-Ambrożej K, *et al.*, 2021) VCI is also commonly used to determine coating integrity and coating thickness in modified-release preparations, in which uniform coating is essential in drug-release control. Furthermore, the method also helps in isolating some flaws like cracks, voids or inconsistencies in layers that may affect the functionality of the products. VCI in sustained-release systems In sustained-release systems using matrices, VCI can be used to understand the spatial distribution of the polymers and APIs, which is used to explain the process of drug release and improve the design of the formulations. Other important benefits are the capability of tracking the changes in solid dosage forms during stability investigations, given that VCI has the capacity of detecting the early indication of degradation, phase separation or recrystallization (Beke ÁK, Gyürkés M, *et al.*, 2021). Moreover, linking VCI and Process Analytical Technology (PAT) allows to monitor the manufacturing process because of blending, granulation, and compression in real-time to maintain the quality of the product. This strategy will be in line with both regulatory requirements of Quality by Design (QbD), which promotes sound process development and validation. In general, Vibrational chemical imaging in solid dosage form characterization provides a multifaceted, non-invasive analytical platform which promotes the knowledge of formulation structure development, provides better quality control and contributes to the creation of more efficient and dependable pharmaceutical products.

Polymorphism and Crystallinity Analysis Using Vibrational Imaging

Crystallinity and polymorphism are very important properties in pharmaceutical development since they play very relevant roles in determining solubility, stability, bioavailability and overall

therapeutic actions of the active pharmaceutical ingredients (APIs). One aspect that has come out as a potent instrument in the characterization and spatial localization of various polymorphic forms and different crystalline states in pharmaceutical systems is vibrational chemical imaging (VCI), especially Fourier Transform Infrared (FTIR), and Raman imaging. These methods take advantage of small differences in molecular vibrations due to differences in crystal lattice structure, and allow the discrimination of polymorphs that could otherwise be indistinguishable using more traditional approaches to evaluation. Specifically, Raman imaging is readily sensitive to crystal structure and is commonly employed in the detection of polymorphic transitions, whereas FTIR imaging is complementary information on the interaction of functional groups and the pattern of hydrogen bonding. VCI enables visualization of the distribution of polymorph in solid dosage forms so that heterogeneity is revealed and it can influence drug dissolution and stability. The interpretation of complex spectral data is usually done through chemometric techniques, including principal component analysis (PCA), multivariate curve resolution (MCR) and cluster analysis to allow the different crystalline forms to be discriminated and quantified. VCI is also used in estimating the extent of crystallinity as well as the existence of amorphous space in formulations, besides polymorphism. Amorphous form of a drug may have greater solubility and a lower physical stability than the crystalline form of a drug and it is thus necessary to measure crystallinity when formulating and storing a drug. (Bunaciu AA, *et al.*, 2024) Vibrational imaging is able to detect amorphous-crystalline transitions which are possible under stress conditions like humidity, temperature, or mechanical processing. Moreover, the method assists in situ monitoring of crystallization reactions, and it gives information regarding the processes of nucleation and growth. The information can be useful in streamlining manufacturing and maintaining uniformity in the quality of its products. VCI is further expanded by the incorporation of both Process Analytical Technology (PAT) and Quality by Design (QbD) frameworks making it even more useful regarding regulatory compliance and quality assurance. Comprehensively, the vibrational chemical imaging will provide a non-destructive and high sensitivity method of analysis of polymorphism and crystallinity, which will help in the formation of stable, effective, and high-quality pharmaceutical products. (Thapa Magar K, *et al.*, 2024)

Drug-Excipient Compatibility and Interaction Studies

The pharmaceutical formulation development necessitates the drug pharmaceutical interaction and the drug-excipient compatibility studies to be carried out because these studies have a direct impact on the

stability, efficacy, and safety of the end dosage form. Vibrational chemical imaging (VCI) and Fourier Transform Infrared (FTIR) especially Raman imaging is an attractive, non-destructive system that is able to study the molecular-scale interactions of active pharmaceutical ingredients (APIs) and excipients in complex formulations. Such methods permit to observe slight alterations of vibrational spectra that occur as a result of chemical interactions, hydrogen bonding, ionic interactions, and van der Waals forces that can modify the physicochemical properties of the drug. FTIR spectroscopy is particularly valuable in revealing how functional groups interrelate and change in their characteristic absorption bands whereas Raman imaging can also be used to provide high spatial resolution in mapping the distribution of interacting components in the dosage matrix. ABA combination with chemometric techniques, including principal component analysis (PCA), multivariate curve resolution (MCR), and cluster analysis, can improve the analysis of multifaceted spectral data, and it is possible to distinguish between reviewable and incompatible systems. By using the methods, VCI is able to detect incompatibilities like degradation reactions or formations of new chemical entities or phase separation which could interfere with product stability. Moreover, the method aids in evaluating the excipient functionality e.g. how it influences the drug release, solubility or stabilizes the amorphous forms. (Martínez LM, *et al.*, 2022) VCI can be used to visualize the distribution of API and excipients and the areas of potential interaction, which can be used to investigate the homogeneity of formulations and microstructural organization. This is of special concern in multi-component systems where local interactions can result into variability of drug action. In addition, VCI facilitates rapid stability tests in that early-stage interactions during stressful conditions like temperature and humidity are identified, thus assisting in projecting stability over time. VCI is used in compatibility studies that comply with regulatory requirements in Quality by Design (QbD), which has ensured that the attributes of materials are well understood and how they influence the quality of the product. Altogether, a combination of vibrational chemical imaging and chemo metric analysis represent a broad-range and sensitive methodology of assessing drug-excipient interactions, which eventually can be used to create stable and robust pharmaceutical formulations.

Quality Control and Regulatory Perspectives

Quality control and regulatory perspectives play a crucial role in the adoption of vibrational chemical imaging (VCI) and chemo metric techniques within the pharmaceutical industry, as they ensure that analytical methods meet stringent standards for accuracy, reliability, and compliance. VCI, encompassing

techniques such as FTIR, Raman, and NIR imaging, provides non-destructive, high-resolution chemical and spatial information that is highly valuable for assessing critical quality attributes (CQAs) of pharmaceutical products, including content uniformity, polymorphic form, crystallinity, and drug–excipient interactions. From a quality control standpoint, these techniques enable rapid and comprehensive analysis of finished dosage forms and raw materials, facilitating early detection of defects, impurities, and inconsistencies that may affect product performance. (Acebedo-Martínez FJ, *et al.*, 2021) The integration of chemo metric models further enhances quality control by enabling robust quantitative and qualitative analysis, supporting batch release testing and ensuring reproducibility across manufacturing processes. Regulatory agencies such as the U.S. Food and Drug Administration, European Medicines Agency, and International Council for Harmonisation emphasize the importance of advanced analytical technologies under frameworks such as Process Analytical Technology (PAT) and Quality by Design (QbD), which promote real-time monitoring, process understanding, and risk-based quality management. VCI aligns well with these

regulatory initiatives by enabling in-line or at-line monitoring of manufacturing processes, such as blending, granulation, and coating, thereby reducing reliance on end-product testing and minimizing batch failures. Furthermore, regulatory acceptance of chemo metric models requires rigorous validation, including assessment of accuracy, precision, specificity, robustness, and model predictability, often following guidelines such as ICH Q2 (R1) for analytical method validation. Data integrity and traceability are also critical considerations, necessitating proper documentation, calibration, and standardization of imaging systems and data processing workflows. Despite its advantages, challenges such as high instrumentation costs, lack of standardized protocols, and the need for skilled expertise can limit widespread implementation. However, ongoing advancements in automation, software integration, and regulatory harmonization are expected to facilitate broader adoption. Overall, the incorporation of VCI and chemometrics into pharmaceutical quality control frameworks enhances analytical capability, supports regulatory compliance, and contributes to the consistent production of safe, effective, and high-quality pharmaceutical products.

Table 1: Comparison of Vibrational Imaging Techniques

Technique	Principle	Advantages	Limitations	Pharmaceutical Applications
FTIR Imaging	Absorption of IR radiation by molecular vibrations	High chemical specificity, fast acquisition (FPA detectors)	Limited spatial resolution, water interference	API distribution, excipient mapping
Raman Imaging	Inelastic scattering of monochromatic light	High spatial resolution, minimal water interference	Fluorescence interference, slower acquisition	Polymorphism, crystallinity, drug mapping
NIR Imaging	Overtone and combination vibrations	Rapid analysis, deep penetration	Lower specificity compared to IR/Raman	Process monitoring, PAT applications
CARS/SRS	Coherent Raman scattering techniques	High sensitivity, rapid imaging	Expensive instrumentation	Real-time imaging, biological systems

Tables 2: Pharmaceutical Applications of VCI and Chemometrics

Application Area	Role of VCI	Role of Chemometrics	Outcome
Solid Dosage Forms	Spatial mapping of API/excipients	Quantitative modeling	Improved content uniformity
Polymorphism Analysis	Identification of crystal forms	Pattern recognition (PCA, MCR)	Stability and bioavailability control
Drug–Excipient Interaction	Detection of molecular interactions	Classification and clustering	Enhanced formulation stability
Quality Control	Non-destructive testing	Predictive modeling (PLS)	Batch consistency
Process Analytical Technology (PAT)	Real-time monitoring	Data integration and automation	Process optimization
Drug Delivery Systems	Visualization of drug distribution	Multivariate analysis	Controlled drug release




Figure1: Overview of Chemometric Methods in Spectral Data Analysis



Figure 2: Quality Control and Regulatory Perspectives

Advances in Imaging Resolution and Data Integration Techniques

Recent developments in imaging resolution and methods to integrate data have greatly increased the possibilities of vibrational chemical imaging (VCI) in pharmaceutical research and studies to allow a more detailed and overall description of complex drug systems. Optical components, detector sensitivity and laser technologies have been improved by great steps and results in significant advances in spatial and spectral resolution, and now chemical heterogeneity can be visualized at micro- scale and even nanoscale. Several

methods have been applied to image at high-resolution above the diffraction limit, including near-field techniques like tip-enhanced Raman spectroscopy (TERS), and confocal Raman microscopy, which is used to study the distribution of drugs, polymorphic domains, and interactions at the interfaces in formulations in detail.(Atre P, *et al.*, 2025) Equally, the development of focal plane array (FPA) sensors in FTIR imaging have enhanced high spectral fidelity of data and high rates of achievement due to their capability to obtain high-throughput data of pharmaceutical samples. Parallel to this, multimodal imaging techniques have been found to

be an effective means of defeating the weaknesses of single techniques. FTIR, Raman, and NIR imaging techniques, when used together in a complementary fashion, the researchers are likely to gain a more comprehensive view of chemical composition and structural organization. Data fusion methods such as low and mid-level and high level fusion allow integration of data sets of various sources to increase accuracy and strength of analysis. In addition, the integration of the state-of-the-art chemometric and machine learning algorithms has transformed the data analysis and made it possible to extract features, classify them, and predict with large hyperspectral datasets automatically. Deep learning and artificial intelligence-related solutions are becoming more and more used in the detection of complex patterns and enhance decision-making in pharmaceutical quality control and process monitoring. Large-scale imaging datasets have also been efficiently handled using cloud-based computing and high-performance data processing platforms, and have enhanced computational efficiency and reproducibility. Such technologies are useful in real-time monitoring and control of Process Analytical Technology (PAT) frameworks, in line with the principles of Quality by Design (QbD). Although these developments have been made, there are still challenges on the standardization of data, its interoperability and approval of integrated models. However, continuous development of imaging resolution and data integration is likely to increase the scope of VCI application to become a pillar of technology in the contemporary pharmaceutical analysis and production.

CONCLUSION

This synergistic approach of high-order spectroscopic imaging, including FTIR, Raman, and NIR imaging with strong multivariate data analysis has made it possible to gain unprecedented understanding on the spatial distribution of, the molecular interaction, and the physicochemical properties of pharmaceutical formulations. These methods overcome significant limitations to traditional methods of analysis, in particular, the impossibility to solve the chemical heterogeneity and spectral overlap in multicomponent systems. With the help of chemometric tools, big and

complicated hyperspectral data can be successfully handled to permit proper qualitative and quantitative data analysis, pattern identification, and predictive modeling. It has found particular application in pharmaceutical uses, in particular in the characterization of solid dosage forms, the determination of polymorphism and crystallinity, in drug-excipient compatibility research, and in quality control. Also, the fact that VCI and chemometrics were aligned with regulatory approaches like Process Analytical Technology (PAT) and Quality by Design (QbD) highlights their significance in the contemporary pharmaceutical production, which allows real-time monitoring, better understanding of the processes, and increased product consistency. Implementation of these technologies will favor the transition of conventional end-product evaluation to a more initiative outlook towards quality assurance using a scientific approach. Imaging resolution, detector technologies, and data integration techniques, such as machine learning and artificial intelligence, have further increased the capabilities of VCI as they can analyze data in high throughput and make decisions automatically. Such innovations not only enhance efficiency in terms of analysis, but also help in production of stronger and more credible pharmaceutical products. In spite of these benefits, there are some weaknesses such as financial instrumentation, special knowledge is required and data standardization and regulatory acceptance of intricate chemometric formulations. To overcome these issues, the interdisciplinary cooperation, technological advancement, and the development of the standardized protocols and the strategies to implement the validation will be needed. In the future, it can be expected that the future of pharmaceutical analysis will be more and more shaped by integrated analytical platforms, which integrate advanced imaging, data science, and real-time process monitoring. The further development of VCI and chemometric methods will be critical in the sustainability of personalized medicine, continuous production, and a next-generation drug delivery system. These newer analytical methods in the end offer a paradigm shift in pharmaceutical sciences by allowing one to gain a deeper insight to the drugs taken by the patients, so as to provide safe, effective and quality medicines to their patients all over the globe.

REFERENCES

- A Gibbons L, Rafferty C, Robinson K, Abad M, Maslanka F, Le N, Raman based chemometric model development for glycation and glycosylation real time monitoring in a manufacturing scale CHO cell bioreactor process. *Biotechnology Progress* 2021; 38.
- Acebedo-Martínez FJ, Alarcón-Payer C, Frontera A, Barbas R, Prohens R, Di Crisci M, Novel Polymorphic Cocrystals of the Non-Steroidal Anti-Inflammatory Drug Niflumic Acid: Expanding the Pharmaceutical Landscape. *Pharmaceutics* 2021; 13: 2140.
- Atre P, Rizvi SAA. Advancements in Pharmaceutical Lyophilization: Integrating QbD, AI, and Novel Formulation Strategies for Next-Generation Biopharmaceuticals. *Biologics* 2025; 5: 35.

- Bakhrushina EO, Shumkova MM, Avdonina YV, Ananian AA, Babazadeh M, Pouya G, Transdermal Drug Delivery Systems: Methods for Enhancing Skin Permeability and Their Evaluation. *Pharmaceutics* 2025; 17: 936.
- Beke ÁK, Gyürkés M, Nagy ZK, Marosi G, Farkas A. Digital twin of low dosage continuous powder blending – Artificial neural networks and residence time distribution models. *European Journal of Pharmaceutics and Biopharmaceutics* 2021; 169: 64–77.
- Bunaciu AA, Hoang VD, Aboul-Enein HY. Fourier transform infrared spectroscopy used in drug excipients compatibility studies. *Applied Spectroscopy Reviews* 2024; 60: 385–403.
- Chen Z, Xue X, Wu H, Gao H, Wang G, Ni G, Visible/near-infrared hyperspectral imaging combined with machine learning for identification of ten *Dalbergia* species. *Front Plant Sci* 2024; 15.
- ChitraRagupathy I, Schweikhard V, Zumbusch A. Multivariate analysis of hyperspectral stimulated Raman scattering microscopy images. *J Raman Spectroscopy* 2021; 52: 1630–1642.
- Dorabiala O, Aravkin AY, Kutz JN. Ensemble Principal Component Analysis. *IEEE Access* 2024; 12: 6663–6671.
- Elhamdaoui O, El Orche A, Cheikh A, Mojemmi B, Nejari R, Bouatia M. Development of Fast Analytical Method for the Detection and Quantification of Honey Adulteration Using Vibrational Spectroscopy and Chemometrics Tools. *Journal of Analytical Methods in Chemistry* 2020; 2020: 1–9.
- Islam MN. Chemometrics in Nondestructive Quality Evaluation, Springer Nature Singapore; 2022, p. 331–355.
- Khadem H, Nemat H, Elliott J, Benaissa M. In Vitro Glucose Measurement from NIR and MIR Spectroscopy: Comprehensive Benchmark of Machine Learning and Filtering Chemometrics. *Heliyon* 2024; 10: e30981.
- Kim EJ, Kim JH, Kim M-S, Jeong SH, Choi DH. Process Analytical Technology Tools for Monitoring Pharmaceutical Unit Operations: A Control Strategy for Continuous Process Verification. *Pharmaceutics* 2021; 13: 919.
- Kucharska-Ambrożej K, Martyna A, Karpińska J, Kiełtyka-Dadasiewicz A, Kubat-Sikorska A. Quality control of mint species based on UV-VIS and FTIR spectral data supported by chemometric tools. *Food Control* 2021; 129: 108228.
- Martínez LM, Cruz-Angeles J, Vázquez-Dávila M, Martínez E, Cabada P, Navarrete-Bernal C, Mechanical Activation by Ball Milling as a Strategy to Prepare Highly Soluble Pharmaceutical Formulations in the Form of Co-Amorphous, Co-Crystals, or Polymorphs. *Pharmaceutics* 2022; 14: 2003.
- Matveeva I, Bratchenko I, Khristoforova Y, Bratchenko L, Moryatov A, Kozlov S, Multivariate Curve Resolution Alternating Least Squares Analysis of In Vivo Skin Raman Spectra. *Sensors* 2022; 22: 9588.
- Mostafa A, Shaaban H. Chemometric Assisted UV-Spectrophotometric Methods Using Multivariate Curve Resolution Alternating Least Squares and Partial Least Squares Regression for Determination of Beta-Antagonists in Formulated Products: Evaluation of the Ecological Impact. *Molecules* 2022; 28: 328.
- Pokhrel DR, Sirisomboon P, Khurnpoon L, Posom J, Saechua W. Comparing Machine Learning and PLSDA Algorithms for Durian Pulp Classification Using Inline NIR Spectra. *Sensors* 2023; 23: 5327.
- Teng G, Wang Q, Cui X, Chen G, Wei K, Xu X, et al. Predictive data clustering of laser-induced breakdown spectroscopy for brain tumor analysis. *Biomed Opt Express* 2021; 12: 4438.
- Thapa Magar K, Boucetta H, Zhao Z, Xu Y, Liu Z, He W. Injectable long-acting formulations (ILAFs) and manufacturing techniques. *Expert Opinion on Drug Delivery* 2024; 21: 881–904.
- Vosough M, Schmidt TC, Renner G. Non-target screening in water analysis: recent trends of data evaluation, quality assurance, and their future perspectives. *Anal Bioanal Chem* 2024; 416: 2125–2136.
- Wang J, Chen Q, Belwal T, Lin X, Luo Z. Insights into chemometric algorithms for quality attributes and hazards detection in foodstuffs using Raman/surface enhanced Raman spectroscopy. *Comp Rev Food Sci Food Safe* 2021; 20: 2476–2507.
- Xie J, Gowen A, Xu W, Xu J. Analysing micro- and nanoplastics with cutting-edge infrared spectroscopy techniques: a critical review. *Anal Methods* 2024; 16: 2177–2197.
- Yang S, Hu X, Zhu J, Zheng B, Bi W, Wang X, Aspects and Implementation of Pharmaceutical Quality by Design from Conceptual Frameworks to Industrial Applications. *Pharmaceutics* 2025; 17: 623.
- Zhu Q, Liu Q, Ma D, Zhu Y, Zhang L, Wang A., Maize Seed Variety Classification Based on Hyperspectral Imaging and a CNN-LSTM Learning Framework. *Agronomy* 2025; 15: 1585.